

THE UNITARITY TRIANGLE: 2002 AND BEYOND

ANDRZEJ J. BURAS

*Technische Universität München, Physik Department,
D-85748 Garching, Germany*

We describe the present status of the Unitarity Triangle and we give an outlook for its future determinations. We discuss new sets of fundamental flavour parameters and comment briefly on new physics beyond the Standard Model.

1 CKM Matrix and the Unitarity Triangle

The unitary CKM matrix^{1,2} connects the *weak eigenstates* (d', s', b') and the corresponding *mass eigenstates* d, s, b :

$$\begin{pmatrix} d' \\ s' \\ b' \end{pmatrix} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix} \equiv \hat{V}_{\text{CKM}} \begin{pmatrix} d \\ s \\ b \end{pmatrix}. \quad (1)$$

Many parametrizations of the CKM matrix have been proposed in the literature. The classification of different parametrizations can be found in³. While the so called standard parametrization⁴ should be recommended⁵ for any numerical analysis, a generalization of the Wolfenstein parametrization⁶ as presented in⁷ is more suitable for my talk. On the one hand it is more transparent than the standard parametrization and on the other hand it satisfies the unitarity of the CKM matrix to higher accuracy than the original parametrization in⁶. Following then the procedure in⁷ we find

$$V_{ud} = 1 - \frac{1}{2}\lambda^2 - \frac{1}{8}\lambda^4, \quad V_{cs} = 1 - \frac{1}{2}\lambda^2 - \frac{1}{8}\lambda^4(1 + 4A^2) \quad (2)$$

$$V_{tb} = 1 - \frac{1}{2}A^2\lambda^4, \quad V_{cd} = -\lambda + \frac{1}{2}A^2\lambda^5[1 - 2(\varrho + i\eta)] \quad (3)$$

$$V_{us} = \lambda + \mathcal{O}(\lambda^7), \quad V_{ub} = A\lambda^3(\varrho - i\eta), \quad V_{cb} = A\lambda^2 + \mathcal{O}(\lambda^8) \quad (4)$$

$$V_{ts} = -A\lambda^2 + \frac{1}{2}A\lambda^4[1 - 2(\varrho + i\eta)] \quad V_{td} = A\lambda^3(1 - \bar{\varrho} - i\bar{\eta}) \quad (5)$$

where

$$\lambda, \quad A, \quad \varrho, \quad \eta \quad (6)$$

are the Wolfenstein parameters with $\lambda \approx 0.22$ being an expansion parameter and terms $\mathcal{O}(\lambda^6)$ and higher order terms have been neglected. A non-vanishing η is responsible for CP violation in

the SM. It plays the role of δ_{CKM} in the standard parametrization. Finally, the bared variables in (5) are given by ⁷

$$\bar{\varrho} = \varrho(1 - \frac{\lambda^2}{2}), \quad \bar{\eta} = \eta(1 - \frac{\lambda^2}{2}). \quad (7)$$

We emphasize that by definition the expression for V_{ub} remains unchanged relative to the original Wolfenstein parametrization and the corrections to V_{us} and V_{cb} appear only at $\mathcal{O}(\lambda^7)$ and $\mathcal{O}(\lambda^8)$, respectively. The advantage of this generalization of the Wolfenstein parametrization over other generalizations found in the literature is the absence of relevant corrections to V_{us} , V_{cd} , V_{ub} and V_{cb} and an elegant change in V_{td} which allows a simple generalization of the unitarity triangle to higher orders in λ as discussed below.

Now, the unitarity of the CKM-matrix implies various relations between its elements. In particular, we have

$$V_{ud}V_{ub}^* + V_{cd}V_{cb}^* + V_{td}V_{tb}^* = 0. \quad (8)$$

Phenomenologically this relation is very interesting as it involves simultaneously the elements V_{ub} , V_{cb} and V_{td} which are under extensive discussion at present. The relation (8) can be represented as a “unitarity” triangle in the complex $(\bar{\varrho}, \bar{\eta})$ plane. One can construct additional five unitarity triangles ⁸ corresponding to other unitarity relations, but I do not have space to discuss them here.

Noting that to an excellent accuracy $V_{cd}V_{cb}^*$ is real with $|V_{cd}V_{cb}^*| = A\lambda^3 + \mathcal{O}(\lambda^7)$ and rescaling all terms in (8) by $A\lambda^3$ we indeed find that the relation (8) can be represented as the triangle in the complex $(\bar{\varrho}, \bar{\eta})$ plane as shown in fig. 1. Let us collect useful formulae related to this triangle:




Figure 1: Unitarity Triangle.

- We can express $\sin(2\phi_i)$, $\phi_i = \alpha, \beta, \gamma$, in terms of $(\bar{\varrho}, \bar{\eta})$. In particular:

$$\sin(2\beta) = \frac{2\bar{\eta}(1 - \bar{\varrho})}{(1 - \bar{\varrho})^2 + \bar{\eta}^2}. \quad (9)$$

- The lengths CA and BA to be denoted by R_b and R_t , respectively, are given by

$$R_b \equiv \frac{|V_{ud}V_{ub}^*|}{|V_{cd}V_{cb}^*|} = \sqrt{\bar{\varrho}^2 + \bar{\eta}^2} = (1 - \frac{\lambda^2}{2}) \frac{1}{\lambda} \left| \frac{V_{ub}}{V_{cb}} \right|, \quad (10)$$

$$R_t \equiv \frac{|V_{td}V_{tb}^*|}{|V_{cd}V_{cb}^*|} = \sqrt{(1 - \bar{\varrho})^2 + \bar{\eta}^2} = \frac{1}{\lambda} \left| \frac{V_{td}}{V_{cb}} \right|. \quad (11)$$

- The angles β and $\gamma = \delta_{\text{CKM}}$ of the unitarity triangle are related directly to the complex phases of the CKM-elements V_{td} and V_{ub} , respectively, through

$$V_{td} = |V_{td}|e^{-i\beta}, \quad V_{ub} = |V_{ub}|e^{-i\gamma}. \quad (12)$$

- The unitarity relation (8) can be rewritten as

$$R_b e^{i\gamma} + R_t e^{-i\beta} = 1. \quad (13)$$

- The angle α can be obtained through the relation

$$\alpha + \beta + \gamma = 180^\circ. \quad (14)$$

Formula (13) shows transparently that the knowledge of (R_t, β) allows to determine (R_b, γ) through

$$R_b = \sqrt{1 + R_t^2 - 2R_t \cos \beta}, \quad \cot \gamma = \frac{1 - R_t \cos \beta}{R_t \sin \beta}. \quad (15)$$

Similarly, (R_t, β) can be expressed through (R_b, γ) :

$$R_t = \sqrt{1 + R_b^2 - 2R_b \cos \gamma}, \quad \cot \beta = \frac{1 - R_b \cos \gamma}{R_b \sin \gamma}. \quad (16)$$

These relations are remarkable. They imply that the knowledge of the coupling V_{td} between t and d quarks allows to deduce the strength of the corresponding coupling V_{ub} between u and b quarks and vice versa.

The triangle depicted in fig. 1, $|V_{us}|$ and $|V_{cb}|$ give the full description of the CKM matrix. Looking at the expressions for R_b and R_t , we observe that within the SM the measurements of four CP *conserving* decays sensitive to $|V_{us}|$, $|V_{ub}|$, $|V_{cb}|$ and $|V_{td}|$ can tell us whether CP violation ($\bar{\eta} \neq 0$) is predicted in the SM. This fact is often used to determine the angles of the unitarity triangle without the study of CP-violating quantities.

2 The Special Role of $|V_{us}|$, $|V_{ub}|$ and $|V_{cb}|$

What do we know about the CKM matrix and the unitarity triangle on the basis of *tree level* decays? Here the semi-leptonic K and B decays play the decisive role. The present situation can be summarized roughly by

$$|V_{us}| = \lambda = 0.221 \pm 0.002 \quad |V_{cb}| = (40.6 \pm 0.8) \cdot 10^{-3}, \quad (17)$$

$$\frac{|V_{ub}|}{|V_{cb}|} = 0.089 \pm 0.008, \quad |V_{ub}| = (3.63 \pm 0.32) \cdot 10^{-3}. \quad (18)$$

implying

$$A = 0.83 \pm 0.02, \quad R_b = 0.39 \pm 0.04. \quad (19)$$

The errors given here look a bit aggressive and should not be considered as giving ranges for the quantities in question. They indicate rather standard deviations. See⁹ for more details. There is an impressive work done by theorists and experimentalists hidden behind these numbers that are in the ball park of various analyses present in the literature. A very incomplete list of references is given in^{10,11}. See also the relevant articles in⁵. Detailed discussions of these analyses with possibly updated values should be available soon¹². In particular the very recent analysis of $|V_{us}|$ ¹³ gives $|V_{us}| = 0.2241 \pm 0.0036$.

The information given above tells us only that the apex A of the unitarity triangle lies in the band shown in fig. 2. While this information appears at first sight to be rather limited, it is very important for the following reason. As $|V_{us}|$, $|V_{cb}|$, $|V_{ub}|$ and consequently R_b are determined here from tree level decays, their values given above are to an excellent accuracy independent of any new physics contributions. They are universal fundamental constants valid in any extension of the SM. Therefore their precise determinations are of utmost importance. In order to answer




Figure 2: “Unitarity Clock”.

the question where the apex A lies on the “unitarity clock” in fig. 2 we have to look at other decays. Most promising in this respect are the so-called “loop induced” decays and transitions and CP-violating B decays. These decays are sensitive to the angles β and γ as well as to the length R_t and measuring only one of these three quantities allows to find the unitarity triangle provided the universal R_b is known.

Of course any pair among (R_t, β, γ) is sufficient to construct the UT without any knowledge of R_b . Yet the special role of R_b among these variables lies in its universality whereas the other three variables are generally sensitive functions of possible new physics contributions. This means that assuming three generation unitarity of the CKM matrix and that the SM is a part of a bigger theory, the apex of the unitarity triangle has to be eventually placed on the unitarity clock with the radius R_b obtained from tree level decays. That is even if using SM expressions for loop induced processes, $(\bar{\rho}, \bar{\eta})$ would be found outside the unitarity clock, the corresponding expressions of the grander theory must include appropriate new contributions so that the apex of the unitarity triangle is shifted back to the band in fig. 2. In the case of CP asymmetries this could be achieved by realizing that the measured angles α , β and γ are not the true angles of the unitarity triangle but sums of the true angles and new complex phases present in extensions of the SM. The better R_b is known, the thinner the band in fig. 2 will be, selecting in this manner efficiently the correct theory. On the other hand as the branching ratios for rare and CP-violating decays depend sensitively on the parameter A , the precise knowledge of $|V_{cb}|$ is also very important.

3 Standard Analysis of the Unitarity Triangle

After these general remarks let us concentrate on the standard analysis of the Unitarity Triangle within the SM. The so-called standard analysis of the UT of fig. 1 involves the values of $|V_{us}|$, $|V_{cb}|$ and $|V_{ub}/V_{cb}|$ extracted from tree level decays, the parameter ε_K that describes the indirect CP violation in $K_L \rightarrow \pi\pi$ decays and the differences of mass eigenstates $\Delta M_{d,s}$ in the $B_d^0 - \bar{B}_d^0$ and $B_s^0 - \bar{B}_s^0$ systems. Setting $\lambda = |V_{us}|$, the analysis proceeds in the following five steps:

Step 1:

From $b \rightarrow c$ transition in inclusive and exclusive leading B-meson decays one finds $|V_{cb}|$ and

consequently the scale of the unitarity triangle:

$$|V_{cb}| \implies \lambda |V_{cb}| = \lambda^3 A. \quad (20)$$

Step 2:

From $b \rightarrow u$ transition in inclusive and exclusive B meson decays one finds $|V_{ub}/V_{cb}|$ and consequently using (10) the side $CA = R_b$ of the unitarity triangle:

$$\left| \frac{V_{ub}}{V_{cb}} \right| \implies R_b = \sqrt{\bar{\varrho}^2 + \bar{\eta}^2} = 4.41 \cdot \left| \frac{V_{ub}}{V_{cb}} \right|. \quad (21)$$

Step 3:

From the experimental value of ε_K and the standard calculation of box diagrams describing $K^0 - \bar{K}^0$ mixing one derives including QCD corrections¹⁴ the constraint $(\lambda = 0.221)$ ¹⁵

$$\bar{\eta} \left[(1 - \bar{\varrho}) A^2 \eta_2 S_0(x_t) + P_c(\varepsilon) \right] A^2 \hat{B}_K = 0.214, \quad (22)$$

where

$$P_c(\varepsilon) = [\eta_3 S_0(x_c, x_t) - \eta_1 x_c] \frac{1}{\lambda^4}, \quad x_i = \frac{m_i^2}{M_W^2}. \quad (23)$$

$S_0(x_t)$ and $S_0(x_t, x_c)$ are known functions^{16,17} and $P_c(\varepsilon) = 0.28 \pm 0.05$ ¹⁸ summarizes the contributions of box diagrams with two charm quark exchanges and the mixed charm-top exchanges. $\hat{B}_K$ is a non-perturbative parameter that represents the relevant hadronic matrix element, the main uncertainty in (22). The short-distance QCD effects are described through the correction factors η_1, η_2, η_3 . The NLO values of η_i with an updated η_1 ¹⁸ are given as follows^{19,20,21}:

$$\eta_1 = 1.45 \pm 0.38, \quad \eta_2 = 0.57 \pm 0.01, \quad \eta_3 = 0.47 \pm 0.04. \quad (24)$$

As illustrated in fig. 3, equation (22) specifies a hyperbola in the $(\bar{\varrho}, \bar{\eta})$ plane. The position of the hyperbola depends on m_t , $|V_{cb}| = A\lambda^2$ and $\hat{B}_K$. With decreasing m_t , $|V_{cb}|$ and $\hat{B}_K$ the ε_K -hyperbola moves away from the origin of the $(\bar{\varrho}, \bar{\eta})$ plane.




Figure 3: Schematic determination of the Unitarity Triangle.

Step 4:

From the observed $B_d^0 - \bar{B}_d^0$ mixing parametrized by ΔM_d and the standard calculation of box diagrams describing this mixing, the side $AB = R_t$ of the unitarity triangle can be determined:

$$R_t = \frac{1}{\lambda} \frac{|V_{td}|}{|V_{cb}|} = 0.86 \cdot \left[\frac{|V_{td}|}{7.8 \cdot 10^{-3}} \right] \left[\frac{0.041}{|V_{cb}|} \right] \quad (25)$$

with

$$|V_{td}| = 7.8 \cdot 10^{-3} \left[\frac{230 \text{ MeV}}{\sqrt{\hat{B}_{B_d} F_{B_d}}} \right] \left[\frac{167 \text{ GeV}}{\bar{m}_t(m_t)} \right]^{0.76} \left[\frac{\Delta M_d}{0.50/\text{ps}} \right]^{0.5} \sqrt{\frac{0.55}{\eta_B}}. \quad (26)$$

Here $\eta_B = 0.55 \pm 0.01$ summarizes the NLO QCD corrections^{20,22} and $F_{B_d} \sqrt{\hat{B}_{B_d}}$ describes the relevant hadronic matrix element. $\bar{m}_t(m_t) = (167 \pm 5) \text{ GeV}$. Note that R_t suffers from additional uncertainty in $|V_{cb}|$, which is absent in the determination of $|V_{td}|$ this way. The constraint in the $(\bar{\varrho}, \bar{\eta})$ plane coming from this step is illustrated in fig. 3.

Step 5:

The measurement of $B_s^0 - \bar{B}_s^0$ mixing parametrized by ΔM_s together with ΔM_d allows to determine R_t in a different manner:

$$R_t = 0.88 \left[\frac{\xi}{1.18} \right] \sqrt{\frac{18.0/\text{ps}}{\Delta M_s}} \sqrt{\frac{\Delta M_d}{0.50/\text{ps}}}, \quad \xi = \frac{\sqrt{\hat{B}_{B_s} F_{B_s}}}{\sqrt{\hat{B}_{B_d} F_{B_d}}}. \quad (27)$$

One should note that m_t and $|V_{cb}|$ dependences have been eliminated this way and that ξ should in principle contain much smaller theoretical uncertainties than the hadronic matrix elements in ΔM_d and ΔM_s separately.

The main uncertainties in this analysis originate in the theoretical uncertainties in the non-perturbative parameters $\hat{B}_K$ and $\sqrt{\hat{B}_d F_{B_d}}$ and to a lesser extent in ξ ²³:

$$\hat{B}_K = 0.86 \pm 0.15, \quad \sqrt{\hat{B}_d F_{B_d}} = (235_{-41}^{+33}) \text{ MeV}, \quad \xi = 1.18_{-0.04}^{+0.13}. \quad (28)$$

The significant uncertainty in ξ is disturbing²⁴ and should be clarified. Also the uncertainty due to $|V_{ub}/V_{cb}|$ in step 2 should certainly be decreased. The QCD sum rules results for the parameters in question are similar and can be found in²⁵. Finally²⁶

$$\Delta M_d = (0.503 \pm 0.006)/\text{ps}, \quad \Delta M_s > 14.4/\text{ps} \text{ at } 95\% \text{ C.L.} \quad (29)$$

4 The Angle β from $B \rightarrow \psi K_S$

One of the highlights of the year 2002 were the considerably improved measurements of $\sin 2\beta$ by means of the time-dependent CP asymmetry in $B_d^0(\bar{B}_d^0) \rightarrow \psi K_S$ decays

$$a_{\psi K_S}(t) \equiv -a_{\psi K_S} \sin(\Delta M_d t) = -\sin 2\beta \sin(\Delta M_d t) \quad (30)$$

The most recent measurements of $a_{\psi K_S}$ from the BaBar²⁷ and Belle²⁸ Collaborations imply

$$(\sin 2\beta)_{\psi K_S} = \begin{cases} 0.741 \pm 0.067 (\text{stat}) \pm 0.033 (\text{syst}) & (\text{BaBar}) \\ 0.719 \pm 0.074 (\text{stat}) \pm 0.035 (\text{syst}) & (\text{Belle}). \end{cases}$$

Combining these results with earlier measurements by CDF ($0.79_{-0.44}^{+0.41}$), ALEPH ($0.84_{-1.04}^{+0.82} \pm 0.16$) and OPAL gives the grand average²⁹

$$(\sin 2\beta)_{\psi K_S} = 0.734 \pm 0.054. \quad (31)$$

This is a mile stone in the field of CP violation and in the tests of the SM as we will see in a moment. Not only violation of this symmetry has been confidently established in the B system, but also its size has been measured very accurately. Moreover in contrast to the constraints of section 3, the determination of the angle β in this manner does not practically suffer from any hadronic uncertainties.

5 Unitarity Triangle 2002

We are now in the position to combine all these constraints in order to construct the unitarity triangle and determine various quantities of interest. In this context the important issue is the error analysis of these formulae, in particular the treatment of theoretical uncertainties. In the literature five different methods are commonly used: Gaussian approach³⁰, Bayesian approach³¹, frequentist approach³², 95% C.L. scan method³³ and the simple (naive) scanning within one standard deviation as used by myself in the past and a few distinguished colleagues of my generation. For the PDG analysis see⁵ and Kleinknechts talk. A critical comparison of these methods will appear soon¹². Recently I have been converted to the Bayesian approach. Consequently, in fig. 4 the result of an analysis in collaboration with Parodi and Stocchi⁹ that uses this approach is shown. The allowed region for $(\bar{\varrho}, \bar{\eta})$ is the area inside the smaller ellipse. We observe that the region $\bar{\varrho} < 0$ is disfavoured by the lower bound on ΔM_s . It is clear from this figure that ΔM_s is a very important ingredient in this analysis and that the measurement of ΔM_s giving R_t through (27) will have a large impact on the plot in fig. 4. Other relatively recent analyses of the UT in the SM can be found in^{30,31,32,34}.




Figure 4: The allowed 95% regions in the $(\bar{\varrho}, \bar{\eta})$ plane in the SM (narrower region) and in the MFV models (broader region) from⁹. The individual 95% regions for the constraint from $\sin 2\beta$, ΔM_s and R_b are also shown. The results are obtained using the fit procedure described in³¹

The ranges for various quantities that result from this analysis are given in the last column of table 1. The first column will be discussed at the end of my talk. The results in this table follow from five steps of section 3 and the direct measurement of $\sin 2\beta$ in (31). They imply in

particular an impressive precision on the angle β :

$$(\sin 2\beta)_{\text{tot}} = 0.725 \pm 0.033, \quad \beta = (23.2 \pm 1.4)^\circ. \quad (32)$$

On the other hand $(\sin 2\beta)_{\text{ind}}$ obtained by using only the five steps of section 3 is found to be⁹

$$(\sin 2\beta)_{\text{ind}} = 0.715^{+0.055}_{-0.045} \quad (33)$$

demonstrating an excellent agreement (see also fig. 4) between the direct measurement in (31) and the standard analysis of the unitarity triangle within the SM. This gives a strong indication that the CKM matrix is very likely the dominant source of CP violation in flavour violating decays. In order to be sure whether this is indeed the case other theoretically clean quantities have to be measured. In particular the angle γ that is more sensitive to new physics contributions than β . In this context the measurement of the ratio $\Delta M_s/\Delta M_d$ will play an important role as for a fixed value of $\sin 2\beta$, the extracted value for γ is a sensitive function of $\Delta M_s/\Delta M_d$.

Strategy	UUT	SM
$\bar{\eta}$	0.369 ± 0.032 (0.298-0.430) [0.260-0.449]	0.357 ± 0.027 (0.305-0.411) [0.288-0.427]
$\bar{\rho}$	0.151 ± 0.057 (0.034-0.277) [-0.023-0.358]	0.173 ± 0.046 (0.076-0.260) [0.043-0.291]
$\sin 2\beta$	$0.725^{+0.038}_{-0.028}$ (0.661-0.792) [0.637-0.809]	$0.725^{+0.035}_{-0.031}$ (0.660-0.789) [0.637-0.807]
$\sin 2\alpha$	0.05 ± 0.31 (-0.62-0.60) [-0.89-0.78]	-0.09 ± 0.25 (-0.54-0.40) [-0.67-0.54]
γ (degrees)	67.5 ± 9.0 (48.2-85.3) [36.5-93.3]	63.5 ± 7.0 (51.0-79.0) [46.4-83.8]
R_b	0.404 ± 0.023 (0.359-0.450) [0.345-0.463]	0.400 ± 0.022 (0.357-0.443) [0.343-0.457]
R_t	0.927 ± 0.061 (0.806-1.048) [0.767-1.086]	0.900 ± 0.050 (0.802-0.998) [0.771-1.029]
ΔM_s (ps^{-1})	$17.3^{+2.2}_{-1.3}$ (15.0-23.0) [11.9-31.9]	$18.0^{+1.7}_{-1.5}$ (15.4-21.7) [14.8-25.9]
$ V_{td} $ (10^{-3})	8.36 ± 0.55 (7.14-9.50) [6.27-10.00]	8.15 ± 0.41 (7.34-8.97) [7.08-9.22]
$ V_{td} / V_{ts} $	0.209 ± 0.014 (0.179-0.238) [0.157-0.252]	0.205 ± 0.011 (0.184-0.227) [0.177-0.233]
$Im\lambda_t$ (10^{-5})	13.5 ± 1.2 (10.9-15.9) [9.4-16.6]	13.04 ± 0.94 (11.2-14.9) [10.6-15.5]

Table 1: Values and errors for different quantities from⁹. In brackets the 95% and 99% probability regions are also given. $\lambda_t = V_{ts}^* V_{td}$.

6 New Set of Fundamental Flavour Variables

During the 1970's and 1980's the variables α_{QED} , the Fermi constant G_F and the sine of the Weinberg angle ($\sin\theta_W$) were the fundamental parameters in terms of which the electroweak tests of the SM have been performed. After the Z^0 boson has been discovered and its mass precisely measured at LEP-I, $\sin\theta_W$ has been replaced by M_Z and the fundamental set used in the electroweak precision studies in the 1990's has been (α_{QED}, G_F, M_Z) . It is to be expected

that when M_W will be measured precisely this set will be changed to (α_{QED}, M_W, M_Z) or (G_F, M_W, M_Z) .

One can anticipate an analogous development in this decade in connection with the CKM matrix. While the set (6) has clearly many virtues and has been used extensively in the literature, one should emphasize that presently no direct independent measurements of η and ϱ are available. η can be measured cleanly in the decay $K_L \rightarrow \pi^0 \nu \bar{\nu}$. On the other hand to our knowledge there does not exist any strategy for a clean independent measurement of ϱ .




Figure 5: The allowed regions (68% and 95%) in the (R_t, β) plane⁹. Different constraints are also shown.

Taking into account the experimental feasibility of various measurements and their theoretical cleanness, the most obvious candidate for the fundamental set in the quark flavour physics for the coming years appears to be⁹

$$|V_{us}|, \quad |V_{cb}|, \quad R_t, \quad \beta \quad (34)$$

with the last two variables measured by means of (27) and (30), respectively. In this context one can investigate, in analogy to the $(\bar{\varrho}, \bar{\eta})$ plane, the (R_t, β) plane for the exhibition of various constraints on the CKM matrix. We show this in fig. 5. Moreover inserting

$$\bar{\varrho} = 1 - R_t \cos \beta, \quad \bar{\eta} = R_t \sin \beta \quad (35)$$

into (2)-(5) and using (13) it is an easy matter to express all elements of the CKM matrix in terms of the variables in (34):

$$V_{ud} = 1 - \frac{1}{2}\lambda^2 - \frac{1}{8}\lambda^4 + \mathcal{O}(\lambda^6), \quad V_{ub} = \frac{\lambda}{1 - \lambda^2/2} |V_{cb}| \left[1 - R_t e^{i\beta} \right], \quad (36)$$

$$V_{cd} = -\lambda + \frac{1}{2}\lambda |V_{cb}|^2 - \lambda |V_{cb}|^2 \left[1 - R_t e^{-i\beta} \right] + \mathcal{O}(\lambda^7), \quad (37)$$

$$V_{cs} = 1 - \frac{1}{2}\lambda^2 - \frac{1}{8}\lambda^4 - \frac{1}{2}|V_{cb}|^2 + \mathcal{O}(\lambda^6), \quad (38)$$

$$V_{tb} = 1 - \frac{1}{2}|V_{cb}|^2 + \mathcal{O}(\lambda^6), \quad V_{td} = \lambda|V_{cb}|R_te^{-i\beta} + \mathcal{O}(\lambda^7), \quad (39)$$

$$V_{ts} = -|V_{cb}| + \frac{1}{2}\lambda^2|V_{cb}| - \lambda^2|V_{cb}|\left[1 - R_te^{-i\beta}\right] + \mathcal{O}(\lambda^6), \quad (40)$$

where in order to simplify the notation we have used λ instead of $|V_{us}|$ as $V_{us} = \lambda + \mathcal{O}(\lambda^7)$.

For the fundamental set of parameters in the quark flavour physics given in (34) we have presently within the SM⁹

$$|V_{us}| = 0.221 \pm 0.002, \quad |V_{cb}| = (40.4 \pm 0.8) \cdot 10^{-3}, \quad R_t = 0.90 \pm 0.05, \quad \beta = (23.2 \pm 1.4)^\circ \quad (41)$$

where the errors represent one standard deviations and the small shift in $|V_{cb}|$ results from the UT fit. The first entry will be soon replaced by 0.2241 ± 0.0036 ¹³.

In the future the situation may change and other sets of fundamental flavour variables could turn out to be more useful than the set (34). As argued in⁹, replacing R_t in (34) by γ could result in the most useful set of flavour variables provided γ can be precisely measured. Similarly the pair (R_b, γ) is very useful as it gives the length of the hand of the unitarity clock in fig. 2 and its position. Other possibilities are discussed in⁹.

7 Outlook: Shopping List

The coming nine years should be very exciting in the field of flavour and CP violation due to a vast amount of data expected from laboratories in Europe, USA and Japan. One should also hope that theorists will sharpen their tools. There are already many reviews of the methods for the extraction of the sides and angles of the UT^{35,36,37,38,39}. Therefore I will be very brief here.

1. It is very desirable that the uncertainties in all inputs entering the five steps of the standard analysis of UT are reduced. The elements $|V_{ub}|$ and $|V_{cb}|$ play here a special role as they are essentially independent of possible new physics contributions. The improved accuracy on ξ in (27) together with a precise measurement of ΔM_s will give us an accurate value of R_t and consequently by means of (15) a prediction for γ . However, the importance of accurate values for $\hat{B}_K$, $F_{B_d}\sqrt{\hat{B}_d}$, and $F_{B_s}\sqrt{\hat{B}_s}$ should not be underestimated. These three quantities are easier to calculate than hadronic matrix elements relevant for non-leptonic K and B decays and are equally important. The precise knowledge of $\hat{B}_K$ combined with improved accuracy on $|V_{cb}|$ will allow to use the precise value of ε_K (step 3) more efficiently. An improved value of $F_{B_d}\sqrt{\hat{B}_d}$ combined with a more accurate value of m_t will give us as seen in (26) a precise value of $|V_{td}|$ and consequently R_t . A precise value of $F_{B_s}\sqrt{\hat{B}_s}$ is important for other reasons. As $|V_{ts}|$ is fixed by the CKM unitarity to be very close to $|V_{cb}|$, the measurement of ΔM_s combined with $F_{B_s}\sqrt{\hat{B}_s}$ allows the measurement of the box diagram function $S_0(x_t)$:

$$S_0(x_t) = 2.32 \left[\frac{270 \text{ MeV}}{\sqrt{\hat{B}_{B_s}} F_{B_s}} \right]^2 \left[\frac{0.040}{|V_{ts}|} \right]^2 \left[\frac{\Delta M_s}{18.0/\text{ps}} \right] \left[\frac{0.55}{\eta_B} \right] \quad (42)$$

and consequently of m_t that could be compared with its direct measurement. This could teach us about the possible new physics beyond SM. For $m_t = 167 \pm 5$ GeV one has $S_0(x_t) = 2.39 \pm 0.12$.

2. The measurement of $\sin 2\beta$ by means of $a_{\psi K_S}(t)$ will certainly be improved in the coming years so that the angle β will be known with an error of 1° ! At this accuracy a closer look at possible theoretical uncertainties will be required. This very precise value for β will be one day

confronted with its value determined by means of clean decays $K_L \rightarrow \pi^0 \nu \bar{\nu}$ and $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ ⁴⁰. With the accuracy for both branching ratios of 10% a measurement of $\sin 2\beta$ with an error ± 0.05 becomes possible. In order to do better not only the accuracy on the branching ratios has to improve but also an NNLO QCD-analysis of these decays combined with improved value of $\bar{m}_c(m_c)$ is required.

In the meantime the CP asymmetry in $B_d \rightarrow \phi K_S$ that also measures $\sin 2\beta$ will be one of the important topics. Being dominated by QCD penguin diagrams it is expected to be more sensitive to new physics than $a_{\psi K_S}(t)$. The first results from BaBar and Belle indicate a value for $\sin 2\beta$ that differs significantly from (31). The recent excitement about this anomaly could be premature, however, as the experimental errors are still large and the decay is not as theoretically clean as $B \rightarrow \psi K_S$. Recent summary is given in ²⁹. See also ⁴¹.

3. Another hot topic is the measurement of $\sin 2\alpha$ through the CP asymmetry in $B_d \rightarrow \pi^+ \pi^-$ that unfortunately is polluted by QCD penguin diagrams and consequently by hadronic uncertainties. There is a vast literature on this subject and many suggestions have been put forward in order to overcome the hadronic uncertainties with the hope to extract the true angle α . Unfortunately the BaBar and Belle data on $a_{CP}(\pi^+ \pi^-)$ disagree with each other with the asymmetry being consistent with zero and large, respectively. Similarly there is no real consensus among theorists. Recent summary is given in ²⁹. See also ⁴². The situation reminds us of ε'/ε at the beginning of the 1990s. Yet, I am convinced that here the experimentalists will reach much faster the agreement than was the case of ε'/ε . Moreover, as the theoretical issues appear to be less involved than in ε'/ε , I expect that some consensus will be reached by theorists in the coming years. On the other hand I have some doubts that a precise value of α will follow in a foreseeable future from this enterprise. However, one should also stress ⁴⁰ that only a moderately precise measurement of $\sin 2\alpha$ can be as useful for the UT as a precise measurement of the angle β . This has been recently reemphasized in ^{43,9}. This is clear from table 1 that shows very large uncertainties in the indirect determination of $\sin 2\alpha$.

4. In view of the comments made in the previous section a precise measurement of the angle γ is of utmost importance. An excellent overview of various strategies for γ can be found in ⁴⁴. The present efforts concentrate around the decays $B_d^0 \rightarrow \pi K$ and $B^\pm \rightarrow \pi K$. On the one hand the data from BaBar and Belle improved considerably this year. On the other hand, there exist several methods like QCDF ⁴⁵ and PQCD ⁴⁶ approaches and more phenomenological approaches: the mixed strategy ⁴⁷, the charged strategy ⁴⁸, the neutral strategy ⁴⁹ and the Wick contraction method ^{50,51}. While I agree to some extent with the Rome group ⁵² that the issue is more involved than stated sometimes by some authors, one cannot deny a great progress made by theorists during the last three years and I am confident that a combination of all $B \rightarrow \pi K$ and $B \rightarrow \pi\pi$ channels will offer in due time a useful, if not the most precise, determination of γ .

Another, very interesting line of attack is to use the U-spin symmetry ^{53,54,55,56} for the determination of γ . In particular the strategies involving the U-spin related decays $B_{d,s}^0 \rightarrow \psi K_S$ and $B_{d,s}^0 \rightarrow D_{d,s}^+ D_{d,s}^-$ ⁵³ and $B_s^0 \rightarrow K^+ K^-$ and $B_d^0 \rightarrow \pi^+ \pi^-$ ⁵⁴ appear to be promising for Run II at FNAL and in particular for LHC-B. They are unaffected by FSI and are only limited by U-spin breaking effects.

Yet, there is no doubt that at the end the most precise determinations of γ will come from the strategies involving $B_d \rightarrow D^{(*)\pm} \pi^\mp$ ⁵⁷ and $B_s \rightarrow D_s^{(*)\pm} K^{(*)\mp}$ ⁵⁸ in which all hadronic uncertainties cancel. One should also mention the triangle construction of Gronau and Wyler ⁵⁹ that uses $B^\pm \rightarrow K^\pm \{D^0, \bar{D}^0, D_\pm^0\}$ where $D_\pm^0$ denotes the CP eigenstates of the neutral D system. However, this method is problematic because of the small branching ratios of the colour suppressed channel $B^+ \rightarrow D^0 K^+$ and its charge conjugate. Variants of this method which could be more promising have been proposed in ^{60,61}.

5. Finally a few rare K and B decays should be put on this shopping list. The recent events for $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ are very encouraging ⁶². In particular one can construct the UT exclusively

by means of $K_L \rightarrow \pi^0 \nu \bar{\nu}$ and $K^+ \rightarrow \pi^+ \nu \bar{\nu}$ ⁴⁰. See also ⁶³. The accuracy of this construction can compete with the one by means of B decays, provided the branching ratios are precisely measured and the uncertainties in $|V_{cb}|$ and m_c reduced. Similarly $K_L \rightarrow \pi^0 \nu \bar{\nu}$ appears to be the best decay to measure the area of the unrescaled unitarity triangle or equivalently $\text{Im}\lambda_t$ in table 1. Finally

$$\frac{Br(B \rightarrow X_d \nu \bar{\nu})}{Br(B \rightarrow X_s \nu \bar{\nu})} = \left| \frac{V_{td}}{V_{ts}} \right|^2, \quad \frac{Br(B_d \rightarrow \mu^+ \mu^-)}{Br(B_s \rightarrow \mu^+ \mu^-)} = \frac{\tau_{B_d} m_{B_d} F_{B_d}^2}{\tau_{B_s} m_{B_s} F_{B_s}^2} \left| \frac{V_{td}}{V_{ts}} \right|^2 \quad (43)$$

allow to determine $|V_{td}|/|V_{ts}|$ or equivalently R_t that can be compared with its determination by means of $\Delta M_d/\Delta M_s$ in (27). As these decays are dominated in the SM by Z^0 -penguin diagrams, while $\Delta M_{d,s}$ are governed by box diagrams, this comparison offers a very good test of the SM.

8 Going Beyond the Standard Model

If the SM is the proper description of flavour and CP violation, all branching ratios and CP asymmetries are given just in terms of four flavour variables, such as the sets (6), (34) or the sets considered in ⁹. This necessarily implies relations between various branching ratios and asymmetries that have to be satisfied independently of the values of the flavour parameters in question if the SM is the whole story. Such relations have been extensively studied in ^{40,65,66,67}.

Now, beyond the SM the amplitude for any decay can be generally written as ⁶⁴

$$A(\text{Decay}) = \sum_i B_i \eta_{\text{QCD}}^i V_{\text{CKM}}^i [F_{\text{SM}}^i + F_{\text{New}}^i] + \sum_k B_k^{\text{New}} [\eta_{\text{QCD}}^k]^{\text{New}} V_{\text{New}}^k [G_{\text{New}}^k]. \quad (44)$$

The non-perturbative parameters B_i represent the hadronic matrix elements of relevant local operators Q_i present in the SM. For instance in the case of $K^0 - \bar{K}^0$ mixing, the matrix element of the operator $\bar{s} \gamma_\mu (1 - \gamma_5) d \otimes \bar{s} \gamma^\mu (1 - \gamma_5) d$ is represented by the parameter $\hat{B}_K$ in (22). There are other non-perturbative parameters in the SM that represent matrix elements of operators Q_i with different colour and Dirac structures.

The objects η_{QCD}^i are the QCD factors analogous to η_i and η_B . Finally, F_{SM}^i stand for the so-called Inami-Lim functions ¹⁶ that result from the calculations of various box and penguin diagrams. They depend on the top-quark mass. An example is the function S_0 in (22).

New physics can contribute to our master formula in two ways. First, it can modify the importance of a given operator, that is relevant already in the SM, through the new short distance functions F_{New}^i that depend on the new parameters in the extensions of the SM like the masses of charginos, squarks, charged Higgs particles and $\tan \beta = v_2/v_1$ in the MSSM. These new particles enter the new box and penguin diagrams. Secondly, in more complicated extensions of the SM new operators (Dirac structures) that are either absent or very strongly suppressed in the SM, can become important. Their contributions are described by the second sum in (44) with $B_k^{\text{New}}, [\eta_{\text{QCD}}^k]^{\text{New}}, V_{\text{New}}^k, G_{\text{New}}^k$ being analogs of the corresponding objects in the first sum of the master formula. The V_{New}^k show explicitly that the second sum describes generally new sources of flavour and CP violation beyond the CKM matrix. This sum may, however, also include contributions governed by the CKM matrix that are strongly suppressed in the SM but become important in some extensions of the SM. A typical example is the enhancement of the operators with Dirac structures $(V - A) \otimes (V + A)$, $(S - P) \otimes (S \pm P)$ and $\sigma_{\mu\nu}(S - P) \otimes \sigma^{\mu\nu}(S - P)$ contributing to $K^0 - \bar{K}^0$ and $B_{d,s}^0 - \bar{B}_{d,s}^0$ mixings in the MSSM with large $\tan \beta$ and in supersymmetric extensions with new flavour violation. The latter may arise from the misalignment of quark and squark mass matrices.

Now, the new functions F_{New}^i and G_{New}^k as well as the factors V_{New}^k may depend on new CP violating phases complicating considerably phenomenological analysis. On the other hand

there exists a class of extensions of the SM in which the second sum in (44) is absent (no new operators) and flavour changing transitions are governed by the CKM matrix. In particular there are no new complex phases beyond the CKM phase. We will call this scenario “Minimal Flavour Violation” (MFV)⁶⁷ being aware of the fact that for some authors MFV means a more general framework in which also new operators can give significant contributions. See for instance the recent discussions in^{68,69}. In the MFV models, as defined here, the master formula (44) simplifies to

$$A(\text{Decay}) = \sum_i B_i \eta_{\text{QCD}}^i V_{\text{CKM}}^i [F_{\text{SM}}^i + F_{\text{New}}^i] \quad (45)$$

with F_{SM}^i and F_{New}^i being real.

Many relations between various quantities valid in the SM are also valid for MFV models or can be straightforwardly generalized to these models. One of the interesting properties of the MFV models is the existence of the universal unitarity triangle (UUT)⁶⁷ that can be constructed from quantities in which all the dependence on new physics cancels out or is negligible like in tree level decays from which $|V_{ub}|$ and $|V_{cb}|$ are extracted. The values of $\bar{\varrho}$, $\bar{\eta}$, α , β , γ , R_b , and R_t resulting from this determination are the “true” values that are universal within the MFV models. Various strategies for the determination of the UUT are discussed in⁶⁷.

The presently available quantities that do not depend on the new physics parameters within the MFV-models and therefore can be used to determine the UUT are R_t from $\Delta M_d/\Delta M_s$ by means of (27), R_b from $|V_{ub}/V_{cb}|$ by means of (10) and $\sin 2\beta$ extracted from the CP asymmetry in $B_d^0 \rightarrow \psi K_S$. Using only these three quantities, we show in figure 4 the allowed universal region for $(\bar{\varrho}, \bar{\eta})$ (the larger ellipse) in the MFV models⁹. The central values, the errors and the 95% (and 99%) C.L. ranges for various quantities of interest related to this UUT are collected in table 1. Similar analysis has been done in⁶⁹.

It should be stressed that any MFV model that is inconsistent with the broader allowed region in figure 4 and the UUT column in table 1 is ruled out. We observe that there is little room for MFV models that in their predictions differ significantly from the SM. It is also clear that to distinguish the SM from the MFV models considered here on the basis of the analysis of the UT only, will require considerable reduction of theoretical uncertainties.

Such a distinction should be much easier in the MSSM with minimal flavour violation but with large $\tan \beta$. Even if the CKM parameters in this model are expected to be very close to the ones in the SM, the presence of neutral Higgs boson penguin-like diagrams with charginos and stop-quarks in the loop can *increase* by orders of magnitude the branching ratios of the rare decays $B_{s,d}^0 \rightarrow \mu^+ \mu^-$ ^{70,71,72,68} and to *decrease* significantly the B_s^0 - $\bar{B}_s^0$ mass difference ΔM_s ⁷³ relative to the expectations based on the SM. Most recent discussions, including also more complicated scenarios, can be found in^{68,73,74,69,75}.

9 Concluding Remarks

The recent direct measurements of $\sin 2\beta$ by BaBar and Belle opened a new era of the precise tests of the flavour structure of the SM and its extensions. These measurements have shown how important it is to have quantities that are free of theoretical uncertainties. With a single direct and clean measurement of the angle β it was possible to achieve accuracy comparable with the indirect measurements of β that involves simultaneously a number of quantities like $|V_{ub}|$, $|V_{cb}|$, $\Delta M_{s,d}$ and ε_K that are all subject to theoretical uncertainties. This lesson makes it clear that one should make all efforts to realize the clean strategies that involve $B_d \rightarrow D^{(*)\pm} \pi^\mp$ and $B_s \rightarrow D_s^{(*)\pm} K^{(*)\mp}$ for γ , $K \rightarrow \pi \nu \bar{\nu}$ for $\sin 2\beta$, $Im \lambda_t$ and the UT as well as the rare decays $B \rightarrow X_{s,d} \nu \bar{\nu}$ and $B_{s,d} \rightarrow \mu^+ \mu^-$ relevant for $|V_{td}|/|V_{ts}|$. Similar comments apply to a number of strategies with small uncertainties as $\Delta M_d/\Delta M_s$ discussed in section 7. However, to this end our theoretical tools have to be improved.

The next years will certainly bring new insight in the flavour structure of the SM and its extensions. In particular it will be important to resolve the issues related to CP asymmetries in $B_d \rightarrow \pi^+\pi^-$ and $B_d \rightarrow \phi K_S$ and to measure ΔM_s . Similarly it will be important to see whether large $\tan\beta$ effects predicted in the MSSM in $B_{s,d} \rightarrow \mu^+\mu^-$ and ΔM_s are realized in nature. As emphasized by Peccei⁷⁶ even more important is the search for new complex phases with the hope to find convincing scenarios that would simultaneously explain the size of CP violation in the low energy processes and the matter-antimatter asymmetry observed in the universe. Another issue is the CP violation in the neutrino sector.

It is conceivable that the physics responsible for the matter-antimatter asymmetry involves only very short distance scales, as the GUT scale or the Planck scale, and the related CP violation is unobservable in the experiments performed by humans. Yet even if such an unfortunate situation is a real possibility, it is unlikely that the single phase in the CKM matrix provides a fully adequate description of CP violation at scales accessible to experiments performed on our planet in this millennium. On the one hand the KM picture of CP violation is so economical that it is hard to believe that it will pass future experimental tests in spite of its recent successes seen in fig. 4. On the other hand almost any extension of the SM contains additional sources of CP violating effects⁷⁷. As some kind of new physics is required in order to understand the patterns of quark and lepton masses and mixings and generally to understand the flavour dynamics, it is very likely that this physics will bring new sources of CP violation modifying KM picture considerably. In any case the flavour physics and CP violation will remain to be a very important and exciting field at least for the next 10-15 years.

Acknowledgments

I would like to thank Fabrizio Parodi and Achille Stocchi for a wonderful collaboration on the matters related to the UT and the organizers for inviting me to such a pleasant conference. I also thank the Max-Planck Institute for Physics in Munich for covering my travel expenses. The work presented here has also been supported in part by the German Bundesministerium für Bildung und Forschung under the contract 05HT1WOA3 and the DFG Project Bu. 706/1-1.

References

1. N. Cabibbo, Phys. Rev. Lett. **10** (1963) 531.
2. M. Kobayashi and K. Maskawa, Prog. Theor. Phys. **49** (1973) 652.
3. H. Fritzsch and Z.Z. Xing, Prog. Part. Nucl. Phys. **45** (2000) 1.
4. L.L. Chau and W.-Y. Keung, Phys. Rev. Lett. **53** (1984) 1802.
5. K. Hagiwara et al., “Review of Particle Physics”, Phys. Rev. **D 66** (2002) 010001.
6. L. Wolfenstein, Phys. Rev. Lett. **51** (1983) 1945.
7. A.J. Buras, M.E. Lautenbacher and G. Ostermaier, Phys. Rev. **D 50** (1994) 3433.
8. R. Aleksan, B. Kayser and D. London, Phys. Rev. Lett. **73** (1994) 18; J.P. Silva and L. Wolfenstein, Phys. Rev. **D 55** (1997) 5331; I.I. Bigi and A.I. Sanda, hep-ph/9909479.
9. A.J. Buras, F. Parodi and A. Stocchi, hep-ph/0207101.
10. Ch.W. Bauer, Z. Ligeti and M. Luke, Phys. Lett. **B479** (2000) 395; Phys. Rev. **D64** (2001) 113004; Th. Becher and M. Neubert, Phys. Lett. **B535** (2002) 127; M. Neubert, Phys. Lett. **B543** (2002) 269; M. Artuso and E. Barberio, hep-ph/0205163; N. Uraltsev, hep-ph/0210044; I.Z. Rothstein, hep-ph/0111337; Ch.W. Bauer, Z. Ligeti, M. Luke and A.V. Manohar, hep-ph/0210027.
11. Results presented at the CKM Unitarity Triangle Workshop, CERN Feb 2002. <http://ckm-workshop.web.cern.ch/ckm-workshop/> in particular see also : LEP Working group on $|V_{cb}|$: <http://lepvcb.web.cern.ch/LEPVcb/> Winter 2002 averages. LEP Working group on

$|V_{ub}|$: <http://battagl.home.cern.ch/battagl/vub/vub.html>.

12. Proceedings of the CKM Workshop at CERN, in progress.
13. V. Cirigliano, G. Colangelo, G. Isidori and G. Lopez-Castro, to appear in¹².
14. A.J. Buras, hep-ph/0101336, lectures at the International Erice School, August, 2000.
15. The numerical value on the r.h.s of (22) being proportional to λ^{-10} is very sensitive to λ but as $A \propto \lambda^{-2}$ and $P_c(\varepsilon) \propto \lambda^{-4}$ the constraint (22) depends only quadratically on λ .
16. T. Inami and C.S. Lim, Progr. Theor. Phys. **65** (1981) 297.
17. A.J. Buras, W. Slominski and H. Steger, Nucl. Phys. **B245** (1984) 369.
18. U. Nierste, recent update.
19. S. Herrlich and U. Nierste, Nucl. Phys. **B419** (1994) 292.
20. A.J. Buras, M. Jamin, and P.H. Weisz, Nucl. Phys. **B347** (1990) 491.
21. S. Herrlich and U. Nierste, Phys. Rev. **D52** (1995) 6505; Nucl. Phys. **B476** (1996) 27.
22. J. Urban, F. Krauss, U. Jentschura and G. Soff, Nucl. Phys. **B523** (1998) 40.
23. L. Lellouch, talk at the ICHEP31, Amsterdam, July 2002, <http://www.chep02.nl>.
24. A.S. Kronfeld and S.M. Ryan, hep-ph/0206058. See also D. Atwood and A. Soni in³⁰.
25. A.A. Penin and M. Steinhauser, Phys. Rev. **D65** (2002) 054006; M. Jamin and B.O. Lange, Phys. Rev. **D65** (2002) 056005; K. Hagiwara, S. Narison and D. Nomura, hep-ph/0205092.
26. LEP Working group on oscillations :
http://lepbose.web.cern.ch/LEPBOSC/combined_results/amsterdam_2002/.
27. B. Aubert et al., BaBar Collaboration, hep-ex/0207042.
28. K. Abe et al., Belle Collaboration, hep-ex/0208025.
29. Y. Nir, hep-ph/0110278.
30. A. Ali and D. London, Eur. Phys. J. **C18** (2001) 665; S. Mele, Phys. Rev. **D59** (1999) 113011; D. Atwood and A. Soni, Phys. Lett. **B508** (2001) 17.
31. M. Ciuchini, G. D'Agostini, E. Franco, V. Lubicz, G. Martinelli, F. Parodi, P. Roudeau, A. Stocchi, JHEP 0107(2001) 013 (hep-ph/0012308).
32. A. Höcker, H. Lacker, S. Laplace, F. Le Diberder, Eur. Phys. J. **C21** (2001) 225.
33. Y. Grossman, Y. Nir, S. Plaszczynski and M.-H. Schune, Nucl. Phys. **B511** (1998) 69; S. Plaszczynski and M.-H. Schune, hep-ph/9911280.
34. M. Bargiotti et al. Riv. Nuovo Cimento **23N3** (2000) 1.
35. The BaBar Physics Book, eds. P. Harrison and H. Quinn, (1998), SLAC report 504.
36. B Decays at the LHC, eds. P. Ball, R. Fleischer, G.F. Tartarelli, P. Vikas and G. Wilkinson, hep-ph/0003238.
37. B Physics at the Tevatron, Run II and Beyond, K. Anikeev et al., hep-ph/0201071.
38. A.J. Buras and R. Fleischer, Adv. Ser. Direct. High. Energy Phys. **15** (1998) 65; hep-ph/9704376.
39. Y. Nir, hep-ph/0109090, lectures at 55th Scottish Univ. Summer School, 2001.
40. G. Buchalla and A.J. Buras, Phys. Rev. **D54** (1996) 6782.
41. R. Fleischer and T. Mannel, hep-ph/0103121; G. Hiller, hep-ph/0207356; A. Datta, hep-ph/0208016; M. Ciuchini and L. Silvestrini, hep-ph/0208087; M. Raidal, hep-ph/0208091.
42. R. Fleischer and J. Matias, hep-ph/0204101; M. Gronau and J.L. Rosner, hep-ph/0202170; D. Atwood and A. Soni, hep-ph/0206045; M. Ciuchini, E. Franco, G. Martinelli and L. Silvestrini, Nucl. Phys. **B 501** (1997) 271.
43. M. Beneke, G. Buchalla, M. Neubert and C.T. Sachrajda, Nucl. Phys. **B 606** (2001) 245; M. Gronau and J.L. Rosner, Phys. Rev. **D65** (2002) 013004; Z. Luo and J.L. Rosner, Phys. Rev. **D54** (2002) 054027.
44. R. Fleischer, hep-ph/0110278, hep-ph/0207324, hep-ph/0208083 and references therein.
45. M. Beneke, G. Buchalla, M. Neubert and C.T. Sachrajda, Phys. Rev. Lett. **83** (1999) 1914, Nucl. Phys. **B 591** (2000) 313, Nucl. Phys. **B606** (2001) 245; G. Buchalla, hep-ph/0202092, M. Beneke, hep-ph/0207228; M. Neubert, hep-ph/0207327; Ch. W. Bauer,

- S. Fleming, D. Pirjol, T.Z. Rothstein and I.W. Stewart, Phys. Rev. **D66** (2002) 014017.
46. H.-Y. Cheng, H.-n. Li and K.-C. Yang Phys. Rev. **D60** (1999) 094005; Y.-Y. Keum, H.-n. Li and A.I. Sanda, Phys. Lett. **B504** (2001) 6, Phys. Rev. **D63** (2001) 054008; H.-n. Li, hep-ph/0101145; A.I. Sanda and K. Ukai, hep-ph/0109001.
 47. R. Fleischer, Phys. Lett. **B365** (1996) 399; R. Fleischer and T. Mannel, Phys. Rev. **D57** (1998) 2752.
 48. M. Neubert and J.L. Rosner, Phys. Lett. **B441** (1998) 403; Phys. Rev. Lett. **81** (1998) 5076.
 49. A.J. Buras and R. Fleischer, Eur. Phys. J. **C11** (1999) 93; Eur. Phys. J. **C16** (2000) 97. For a recent numerical analysis see M. Bargiotti et al., hep-ph/0204029 and the first paper in ⁴².
 50. M. Ciuchini, E. Franco, G. Martinelli and L. Silvestrini, Nucl. Phys. **B501** (1997) 271.
 51. A.J. Buras and L. Silvestrini, Nucl. Phys. **B569** (2000) 3.
 52. M. Ciuchini, E. Franco, G. Martinelli, M. Pierini and L. Silvestrini, Phys. Lett. **515** (2001) 33 and hep-ph/0208048.
 53. R. Fleischer, Eur. Phys. J. **C10** (1999) 299, Phys. Rev. **D60** (1999) 073008.
 54. R. Fleischer, Phys. Lett. **B459** (1999) 306.
 55. M. Gronau and J.L. Rosner, Phys. Lett. **B482** (2000) 71; C.W. Chiang and L. Wolfenstein, Phys. Lett. **B493** (2000) 73.
 56. M. Gronau, Phys. Lett. **B492** (2000) 297.
 57. R.G. Sachs, EFI-85-22 (unpublished); I. Dunietz and R.G. Sachs, Phys. Rev. **D37** (1988) 3186 [E: Phys. Rev. **D39** (1988) 3515]; I. Dunietz, Phys. Lett. **427** (1998) 179; M. Diehl and G. Hiller, Phys. Lett. **517** (2001) 125;
 58. R. Aleksan, I. Dunietz and B. Kayser, Z. Phys. **C54** (1992) 653; R. Fleischer and I. Dunietz, Phys. Lett. **B387** (1996) 361. A.F. Falk and A.A. Petrov, Phys. Rev. Lett. **85** (2000) 252; D. London, N. Sinha and R. Sinha, Phys. Rev. Lett. **85** (2000) 1807.
 59. M. Gronau and D. Wyler, Phys. Lett. **B265** (1991) 172.
 60. M. Gronau and D. London, Phys. Lett. **B253** (1991) 483; I. Dunietz, Phys. Lett. **B270** (1991) 75.
 61. D. Atwood, I. Dunietz and A. Soni, Phys. Rev. Lett. **B78** (1997) 3257. M. Gronau, Phys. Rev. **D58** (1998) 037301.
 62. S. Adler et al., (AGS E787) Phys. Rev. Lett. **79** (1997) 2204, Phys. Rev. Lett. **84** (2000) 3768, hep-ex/0111091.
 63. A.J. Buras, Phys. Lett. **333** (1994) 476; Y. Grossman and Y. Nir, Phys. Lett. **398** (1997) 163; G.D'Ambrosio and G. Isidori, Phys. Lett. **530** (2002) 108.
 64. A.J. Buras, hep-ph/0109197, talk given at Kaon 2001, Pisa, June, 2001.
 65. A.J. Buras and R. Fleischer, Phys. Rev. **D64** (2001) 115010.
 66. A.J. Buras and R. Buras, Phys. Lett. **501** (2001) 223; S. Bergmann and G. Perez, Phys. Rev. **D64** (2001) 115009, JHEP **0008** (2000) 034. S. Laplace, Z. Ligeti, Y. Nir and G. Perez, Phys. Rev. **D65** (2002) 094040.
 67. A.J. Buras, P. Gambino, M. Gorbahn, S. Jäger and L. Silvestrini, Phys. Lett. **B500** (2001) 161.
 68. C. Bobeth, T. Ewerth, F. Krüger and J. Urban, Phys. Rev. **D64** (2001) 074014; hep-ph/0204225. J. Urban, hep-ph/0210286.
 69. G. D'Ambrosio, G.F. Giudice, G. Isidori and A. Strumia, hep-ph/0207036.
 70. K.S. Babu and C. Kolda, Phys. Rev. Lett. **84** (2000) 228.
 71. P.H. Chankowski and L. Ślawniowska, Phys. Rev. **D63** (2001) 054012; Acta Phys. Polon. **B32** (2001) 1895.
 72. C.-S. Huang, W. Liao, Q.-S. Yan and S.-H. Zhu, Phys. Rev. **D63** (2001) 114021, Erratum: Phys. Rev. **D64** (2001) 059902.

- 73. A.J. Buras, P.H. Chankowski, J. Rosiek and L. Ślawnowska, Nucl. Phys. **B619** (2001) 434, hep-ph/0207241, hep-ph/0210145.
- 74. G. Isidori and A. Retico, JHEP **0111** (2001) 001 (hep-ph/0110121), hep-ph/0208159.
- 75. A. Dedes and A. Pilaftsis, hep-ph/0209306.
- 76. R.D Peccei, hep-ph/0209245.
- 77. A. Masiero and O. Vives, hep-ph/0104027; P.H. Chankowski and J. Rosiek, Acta Phys. Polon. **B33** (2002), 2329 (hep-ph/0207242); L. Silvestrini, hep-ph/0210031 and refs. therein.